

RESEARCH ARTICLE

LITHO-STRUCTURAL MAPPING USING REMOTE SENSING AND FIELD WORK TECHNIQUES: A CASE STUDY FROM CENTRAL SALT RANGE, DISTRICT KHUSHAB PUNJAB PAKISTAN

Muhib Ullah Khan^{a*}, Touseef Ahmad Khan^a, Umer Sadiq Khan^b, Naveed Ur Rahman^c

^aInstitute of Mountain Hazards and Environment, Chinese Academy of Sciences, Chengdu, 610041, China.

^bAssistant Professor, Department of Computer Science Mir Chakar Khan Rind University Sibi.

^cCollege of Geosciences, China University of Petroleum Beijing, 102200, China.

*Corresponding Author Email: muhibkpk@yahoo.com

This is an open access journal distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

ARTICLE DETAILS

ABSTRACT

Article History:

Received 19 March 2022

Accepted 21 April 2022

Available online 31 May 2022

Litho-structural mapping in an inaccessible rugged and mountainous region like the Central Salt range has frequently been a great challenge and importance for geologists. For this purpose, in the present study, litho-structural mapping was carried out based on enhancement and interpretation of Landsat 8 USGS data by the application of remote sensing and GIS technology and further verified by the data which was collected in field work to the study area. Different band combinations of the image are applied for visual image interpretation to digitize the rock boundaries while as Sobel and Laplacian filtering techniques are utilized for lineament mapping. The purpose of our field work was a comprehensive mapping of Geology and related structural aspects of Kufri, Jhalar, Channaki, Surraki, Khura and Naushehra area, Central Salt Range, district Khushab Punjab Pakistan in the Central Salt Range Pakistan. It includes Litho-structural Mapping at scale of 1:10,000. The results and conclusion of this study demonstrate that the processing and interpretation of Satellite data set can be employed as a powerful tool to improve lithological discrimination and enhance the overall mapping performance in the rugged and hilly terrain like Salt range.

KEYWORDS

Mapping; Remote sensing; GIS; Salt range; Digitization; Field data.

1. INTRODUCTION

Lithological mapping is a very important component of geology. These Lithological maps are very significant in distributing rocks of different types over the earth surface. The history of lithological maps dates back to late 18th and 19th centuries, since then contributed significantly when it comes to understanding the earth's history and help save many problems related to land use such as frequently occurring earthquakes, volcanic activities, underground water quality control and many other natural hazards like flooding and landslides. In the past Litho-structural mapping was carried out by geologists traveling for miles through tough terrains observing the characteristics of rocks and marking different structures and details for that area based on those observed information's. These lithological maps provide huge amount of information. The colors used in lithological maps as patterns mark the distribution of different geologic features on earth surface.

Recent advancements in the available techniques includes interpretation from aerial photography to satellites based remote sensing. As long as the advancement in the field of remote sensing and GIS continues researchers have achieved great success related to different aspects of geosciences and has become as a first-hand tool in lithological mapping. Researchers are very frequently using data from satellites and remote sensing techniques for identification of various lithology-based research. Previous research deduced different approaches discriminate and map different

rock types using multispectral data like principal component analysis (PCA) data, band ratio data, classification based on multiband data, etc (Davis et al., 1993; Gillespie et al., 1986; and Pirasteh et al., 2007).

Salt range is composed of a very complex geology associated with active Himalayan terrain which makes the mapping in region a difficult task to carry out, but satellites composites have proven very useful for the identification and demarcation of these complex geologic segments Remote sensing technology based on multi spectral satellites can efficiently help build lithological maps of those geologically complex regions of Himalayas at very low cost. In order to confirm the accuracy and precision of the obtained results from these digital based techniques requires further information through field work. This present research is carried out to enhance the quality of lithological mapping. This paper first introduces the location and tectonics of the study area and then discuss the methodology of the paper. After that it discuss the procedure of field data and digitization of different lithological units. At the end it discusses result and conclusion of the research.

2. LOCATION AND ACCESSIBILITY OF STUDY AREA.

The study area soon valley is in the north west of Khushab District, Punjab, Pakistan. Its largest settlement is the town of Noshera. The valley extends from the village of Padhrar to Sakesar, the highest peak in the Salt Range. The valley is 35 miles (56 km) long and has an average width of 9 miles (14 km). It covers a 300-square-mile (780 km²) area.

Quick Response Code



Access this article online

Website:

www.earthsciencesmalaysia.com

DOI:

10.26480/esmy.02.2022.81.85




Figure 4: (a) Contour map and (b) 3D surface map of the study area prepared for field work by surfer 14.

4.2 Field Work Data

Before proceeding to the field for ground checks, proper planning has to be made regarding the type of data to be collected and the locations to be selected for this purpose based on the road network. By using Topographic map (figure 3) of the study area, road map of the study area was prepared in order to reach the places to cross check the lithology and lineaments on image with lithology and structural features in the field. Almost ten sites were chosen for the verification of digitized lithology from satellite image. After doing all in the lab, results obtained were checked in the field. For the field work, we have taken with us the digitized map of lithology and lineament of Soon Sakesar valley and the also the map showing the places where we have to collect data related to different rock types and structures. So, ground data form an important source of information for mapping and accuracy estimation.




Figure 5: (a) Mianwali formation Kathwai member near Surakki village (b) Intra formational conglomerate of Kamlial Formation at the south of kufri village (c) Solution holes in Sakesar Limestone At the south of Mustafabad village (d) Hangu Formation Laterite bands at Rakh Surakki village.

4.3 Digitization of Lithological Units

Special software's were used for the purpose to digitization so as to help delineate the spatial extent for those features which we are investigating. The process of digitization mainly involves the conversion of lithological unit's contacts and highlighting lineaments through imagery (based on tones colors, textures, shape and different patterns) and converting the existing geological map to digital formats using Arc GIS (Ali and Ali, 2013). Enhancement of the image from 421 band combinations and making it more contrasting can have good results while performing lithological mapping. Lithologies marked on geological maps are in close relation to the types of rocks marked and indicated on satellite images which were further verified by field work. Identification and digitalization of the lineaments were carried out through satellite images with respect to correlation with roads and the pattern of drainage in the area. These

digitized rock types from Landsat 8 USGS data images were also verified carrying out field in the area and after the verification a complete and precise final map was deduced for the area which is showing in the results portion of the paper.




Figure 6: Landsat 8(USGS) Images Which Has Been Used Further in Digitization and Mapping for Field Work

5. RESULTS AND DISCUSSIONS

Different kind of techniques were applied to enhance Land sat 8 data images (figure 6) which helps converting lithological units and their contacts into a format that is more digital and modern. The main hinderance while identifying the lithology was caused by the vegetation cover. These vegetational covers make it difficult to capture direct radiations which are very important in any lithological interpretation work. Composites of different colors must be created to display an image having multi spectral in color i.e Green, Red and Blue every one of which exhibits its own characteristics. Combining different colors patterns are tested and found useful in distinguishing various lithological boundaries hence selected while digitizing different lithological units using visual image interpretations. Different rock Lithologies identified with the help of data from satellite differ from each other in their texture, tone, drainage pattern system. The results are also confirmed in field. The list of observed formations and their description are given in Table 1.

Table 1: List of Observed Formations

Age	Formation	Lithology	Description
Pleistocene	Kalabagh Conglomerate		Conglomerate
Miocene	Kamlial Formation		Sandstone
Eocene	Sakesar Formation		Limestone and marl
	Nammal Formation		Limestone and shale
Paleocene	Patala Formation		Shale
	Hangu Formation		Sandstone and laterite
Triassic	Mianwali Formation		Limestone, shale and dolomite
Permian	Wargal Limestone		Limestone
	Warcha Sandstone		Sandstone and shale interbeds
	Dandot Formation		Sandstone and shale interbeds
	Tobra Formation		Tillitic Facies, Fresh Water Facies

LEGEND			
Lithology: Conglomerate	Lithology: Marl	Lithology: Laterite	Lithology: Dolomite
Lithology: Limestone	Lithology: Shale	Lithology: Sandstone	Lithology: Tillite

Another aspect of the study includes drainage pattern which is use as additional aid during lithological mapping. Examining the patterns indicated by drainage or drainage density can provide handful clues about lithology even if the rocks under the study are lacking direct exposure. The reason for this can be more, drainage density of hard rocks(basalts) as it is difficult for water to infiltrate down compared to rocks having loss textures (alluvium) where infiltration of water downward is easy which results in low drainage density.

5.1 Mapping & Structure Set Up of the Study Area

The basic purpose of field work was geological mapping and verification of the GIS based data. Toposheet constitutes the base map for geological mapping. Toposheet number 43 D/2 of the Geological Survey of Pakistan was enlarged up to 5 times for the mapping purpose. The project area is located in Central Salt Range, where the zone of decollement has been provided by the evaporites of Salt Range Formation. All the structures are controlled by Salt Range Thrust. Normal faulting is common in the Study area and drag folding and many structures associated with the normal faulting are obvious in the study area. Ridges and depressions area also common in the study area. Ridges are formed of younger formations while depressions are formed of older formations filled with recent to sub recent alluvium.

5.1.1 Dhok Patial Thrust

A section line E-F is drawn in N-S direction. A thrust fault observed along this cross section named as Dhok Patial thrust. Wargal Limestone is overriding the Sakesar Limestone, are exposed along its North-Western flank and South-Eastern flank. Also, section line M-N is drawn in N-S direction. A thrust fault is observed along this cross section named as Dhok Patial Thrust. Chhidru formation is overriding the Minawali formation, are exposed along its North Western and South-Eastern flank.

5.1.2 Dhok Patial Syncline

A section line M-N is drawn in N-S direction. A syncline is observed along this cross section having Alluvium in its core and Sakesar Limestone, Hangu Formation, Minawali Formation and Chhidru formation in its eastern and southern flank.

5.1.3 Sukh Wahan Thrust

A section line E-F is drawn in N-S direction. A thrust fault observed along this cross section named as Sukh Wahan thrust. Chhidru Formation is overriding the Mianwali Formation, are exposed along its North-Western flank and South-Eastern flank. Also, section line G-H is drawn in NW-SE direction. A thrust is observed along this cross section near Sukh Wahan, named as Sukh Wahan Thrust. Chhidru formation is overriding the Minawali formation near Sukh Wahan observed in this section.

5.1.4 Sukh Wahan Anticline

A section line E-F is drawn in N-S direction. An anticline observed along this cross section named as Sukh Wahan anticline having Wargal limestone in its core. Mianwali Formation and Hangu Formation are exposed along its North-Western flank and South-Eastern flank.

5.1.5 Sukh Wahan Syncline

A section line E-F is drawn in N-S direction. A Syncline observed along this cross section near Sukh Wahan, named as Sukh Wahan Syncline having kalabagh conglomerate in its core and Sakesar Limestone, Hangu Formation, Minawali Formation and Chhidru formation are present along its eastern and southern flank.

5.1.6 Jhalar Thrust

A section line M-N is drawn in N-S direction. A thrust is observed along this cross section near Jhalar, named as Jhalar Thrust. Minawali Formation is overriding the Sakesar lime Stone near Jhalar and observed in the section.

5.1.7 KHURA ANTICLINE

A section line G-H is drawn in NW-SE direction. An Anticline is observed along this cross section near Khura, named as Khura Anticline having Wargal limestone in its core and Minawali formation at the top is eroded.

5.1.8 Khura Normal Fault

A section line G-H is drawn in NW-SE direction. A Normal fault is observed along this cross section near Khura, named as Khura Normal fault where Chhidru formation is overriding the Minawali formation.

5.1.9 Sodhi Thrust

A section line B-C is drawn in N-S direction. A thrust is observed along this cross section near Sodhi, named as Sodhi Thrust. Minawali formation is overriding the Sakesar Limestone near Sodhi observed in this section.

5.1.10 Chamnaki Anticline

A section line B-C is drawn in N-S direction. An anticline observed along this cross section near Chamnaki, named as Chamnaki anticline having Wargal limestone in its core. Minawali formation and Chhidru formation are present along its Southern and Eastern flank and at the top Minawali formation is eroded. Also observed in section line G-H.

5.1.11 Chamnaki Syncline

A section line B-C is drawn in N-S direction. A Syncline observed along this cross section near Chamnaki, named as Chamnaki Syncline having Alluvium in its core and Sakesar Limestone, Hangu Formation, Minawali Formation and Chhidru formation are present along its eastern and southern flank. And section line G-H is drawn in NW-SE direction. A syncline is observed having Sakesar Limestone in its Core and Minawali Formation and Chhidru formation in its eastern and southern flank. Also observed in section line M-N is drawn in N-S direction.

5.1.12 Chamnaki Thrust

A section line B-C is drawn in N-S direction. A Thrust observed along this cross section near Chamnaki, named as Chamnaki Thrust. Chhidru formation is overriding the Minawali formation.




Figure 7: Cross Section along the line A-B, B-C, E-F, G-H, M-N.

While mapping lineaments, lithology and drainage network in the study area guided us to the conclusion that the study area has undergone immense deformation That signifies the influence of a major tectonic structure that surrounds the Central parts of Salt Range (Abied, 2008 and Abubaker, 2015). Hence, the present research demonstrates the importance of remote sensing and GIS to identify lithologies and lineaments in those area that are geologically very active and form complex terrains like the one we have in Central Salt Range.




Figure 8: Litho-structural Map of Kufri, Jhalar, Chamnaki, Surakki, Khura And Naushehra Area, Central Salt Range, District Khushab Punjab Pakistan.

6. CONCLUSION

Techniques of remote Sensing and GIS are integrated so that to distinguish different rocks types and lineaments. The high-resolution data used for these integrated interpretations. Functions like GIS spatial analysis helps develop lithological maps with the use of different information layers that were generated, use techniques based on image processing like Principle component analysis, visual image interpretation, filtering and FCC. The type of rocks marked with the help of data from Satellite indicates a very close resemblance to those rocks present in already existing conventional maps including limestone, Conglomerate, basalt, granite, Phyllites and alluvium. The technique of edge enhancement has shown good accuracy and results for better recognition of lineaments that are tough to trace down in field mapping. Although there are some hinders when we use the techniques of satellite remote sensing in the pursue of carrying out lithological mapping as it will indicate some sort of spectral similarity among some lithological units maybe be due to same kind of vegetational covers but we can remove these difficulties by field work. Further it is very hard to identify lineaments on ground but can be very easy to see from Satellite imaging. Hence, Satellite imaging can be very useful to enhance the quality of lithological mapping and identification of lineaments.

REFERENCES

- Abied, A.E., 2008. GIS and remote sensing technology in geological mapping using GIS and remote sensing application in Libya. In Proceedings first conference. Tripoli: Office of Scientific Research and Development.
- Ali, S. A., Ali, U., 2015. Litho-Structural Mapping of Sind Catchment (Kashmir Basin), NW Himalaya, using Remote Sensing & GIS Techniques. *Int. j. sci. res. (IJSR)*, 4 (7), Pp. 1325-1330.
- Butler, R.W.G., Coward, M.P., Harwood, G.M., Knipe, R.J., 1987. Salt, its control on thrust geometry, structural style and gravitational collapse along the Himalayan Mountain front in the Salt Range of northern Pakistan. *Dyna. geo. of salt. Rel. str. Acad. press, Orlando, Florida*, Pp. 339-418.
- Coward, M. P., Butler, R.W.H., 1985. Thrust tectonics and the deep structure of the Pakistan Himalaya. *Geology*, 13 (6), Pp. 417 - 420.
- Davis, P.A., Breed, C.S., McCauley, J.F., Schaber, G.G., 1993. Surficial geology of the Safsaf region, south-central Egypt, derived from remote sensing and field data. *R. sens. environment.*, 46, Pp. 183-203.
- Davis, D.M., Terry, E., 1985. The role of salt in fold-and-thrust belts. *Tectonophysics*, 119 (1-4), Pp. 67-88.
- Ezabti, A., Jovanovic, V., 2015. GIS and remote sensing application in geological mapping and 3D terrain modeling: a case study in eghei uplift, Libya. In proceeding SYNTHESIS Int. sci. conf. IT. Bus. Rel. res. Belgrade, Singidunum University, Serbia, Pp. 615-619.
- Gillespie, A.G., Kahle, A.B., Walker, R.E., 1986. Color enhancement of highly correlated images. I. Decorrelation and HSI contrast stretches. *R. sens. environment.*, 20, Pp. 209-235.
- Lillie, R.J., Johnson, G.D., Yousaf, M., Sher, A.H.Z., Robert, S., 1987. Structural development within the Himalayan foreland fold and thrust belt of Pakistan. *Sed. bas. bas. form. mech. — memoir.*, 12, Pp. 379-392.
- Pirasteh, S., Ali, S.A., Hussaini, H.S., 2007. Morphometric and Structural Analysis in Zagros Mountains, Southwest Iran—An Application of Geo-Information Technology. *J. geom.*, 1 (2), Pp. 39-44.
- Qayyum, M., 1991. Crustal Shortening and Tectonic Evolution of the Salt Range in Northwest Himalayas, Pakistan. Master's Thesis, Oregon State University, Corvallis, Oregon, Pp. 1-124.
- Sameeni, S.J., 1997. Biostratigraphy of the Eocene succession of the Salt Range, Northern Pakistan. Ph.D. Thesis, Punjab University, Lahore, Pp. 64.
- Sarwar, G., De Jong, K.A., 1979. Arcs, oroclines, syntaxes: the curvatures of mountain belts in Pakistan. *Geodyn. of Pakistan. Geo. surv. of Pakistan*, Pp. 341-350.
- Seeber, L., Armbruster, J.G., 1981. Great detachment earthquakes along the Himalayan arc and long-term forecasting. In *Earthquake Prediction – An International Review. Maurice Ewing Series*, 4, Pp. 259-277.
- Yeats, R.S., Lawrence, R.D., 1984. Tectonics of the Himalayan thrust belt in northern Pakistan. *Mar. geo. and oceanogr. of arab. sea and coast. Pakistan*, Pp. 177-198.

