

RESEARCH ARTICLE

PETROLOGICAL AND MECHANICAL PROPERTIES OF THE DAHOMEYAN GRANITIC ROCKS – A CASE STUDY AT GREEN VALLEY AND AKROFU

Matthew Coffie Wilson^{a*}, Emmanuel Kwesi Arkaifie^b, Bernard Cudjoe^c, Jeremiah Mensah^d, Bernard Audinada Ampofo^a

^a Kwame Nkrumah University of Science and Technology, Department of Geological Engineering, University Post Office, Kumasi–Ghana.

^b Spectech Engineering Services Limited, Plot 9, Helena Junction, Airport Ridge Residential Area. PMB Ax 20, Takoradi – Ghana.

^c Cardinal Resources Limited, Durugu Residential Area, Kumbosco, Bolgatanga – Ghana.

^d Community Water and Sanitation Agency, Tarkwa-Nsuaem Municipal Assembly. Post Office Box 1, Tarkwa – Ghana.

*Corresponding Author Email: regimatt2003@yahoo.co.uk; mcwilson.coe@knust.edu.gh

This is an open access article distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ARTICLE DETAILS

Article History:

Received 26 June 2022

Accepted 28 July 2022

Available online 01 August 2022

ABSTRACT

The main aim of this project is to determine the structures, petrography, strength and competence of Dahomeyan granitic rocks at Green Valley and Akrofu in Ghana and compare them from both localities in terms of their mineralogical composition and strength properties. Thin sections were prepared with rock samples from the field at the KNUST Geological Engineering Laboratory and petrographic microscope used to determine the different types of minerals in the samples and also the rock type. The mechanical strengths of the rocks were determined at the KNUST Civil Engineering Laboratory using Schmidt Hammer device. The studied Dahomeyan rocks at both Akrofu and Green Valley are respectively classified as granodiorite and biotite granitic gneiss. The mineralogical composition of different modal percentages includes quartz, feldspars, biotite, muscovite and hornblende. The presence of fractures and foliations due to a mafic biotite mineral indicate deformations rocks. The granodiorite is classified as weak rock with compressive strength of 23.50 MPa. This is due to higher fractures in the rock. In conclusion, the higher the presence of high grade of fractures and foliations due to biotite mineral, the weaker the strength of the rock. The biotite granite gneiss at Green Valley can thus be useful as crushed stone in road construction as well as for building material. For the rock texture, coarser mineral fabric corresponds to higher uniaxial compressive strength. The strength of these granitoids were influenced by the climatological changes as well as the intensity of weathering they had undergone within the area.

KEYWORDS

Dahomeyan granitic rocks, petrography, strength properties, granitic rocks, rock structures

1. INTRODUCTION

The Dahomeyan rock group is Ghana's easternmost rock group, and it differs significantly from other Ghanaian rocks in that it is composed of high-grade metamorphic rocks. It is located in the southeastern corner of Ghana, roughly along a line drawn north-northeast from Accra to intersect the Ghana-Togo border near Agome in Togo. The system, which lies beneath the Accra planes, covers an area of approximately 7000 square kilometers. The Dahomeyan System is made up of Birimian or younger formations that were involved in the Pan African thermo-tectonic event (Kennedy, 1964). The system is made up of four granitic and mafic gneiss lithologic belts. Mafic gneisses are igneous oligoclase, andesine, hornblende, salite, and garnet gneisses with relatively uniform oligoclase, andesine, hornblende, salite, and garnet compositions (Holm, 1974).

Petrography is the detailed study of rocks; the description in the field, macroscopic and microscopic analysis of minerals under a petrographic microscope. Mechanical Properties are physical properties that a material exhibits upon the application of forces. Most of the previous studies conducted in the Dahomeyide basement terrane focused on the orthogneisses without considering the significance of the granites (Attoh

et al., 1997). This study highlights on providing detailed information on the structural features of the minerals as well as highlighting on the petrographic and mechanical properties of the Dahomeyan granitic rocks from the Green Valley and Akrofu localities. The results obtained from this project work will be very vital in knowing the right type of rock and its competences to choose when performing any engineering works. The results obtained from this project work will be useful for future reference and used in engineering investigations. The main aim of this project is to determine the petrographic and mechanical properties of the Dahomeyan granitic rocks in Green Valley and Akrofu. The research helps to determine the mineralogical composition of the rocks; determine the strength and competence of the rocks; and compare rocks from both localities in terms of their mineralogical composition and strength properties.

2. GEOLOGIC SETTINGS

The Pan African orogenic Dahomeyide belt formed as a result of the continental collision between the eastern margin of the West African Craton (WAC) and the Benin-Nigerian shield (Figure 1a) after the closure of a subduction event around 0.6 Ga years ago (Caby, 1989; Affaton et al., 1991; Ganade de Araujo et al., 2016). The belt has a well-organized

Quick Response Code



Access this article online

Website:

www.earthsciencesmalaysia.com

DOI:

10.26480/esmy.02.2022.114.119

4.1.2 Microscopic Analysis

The thin sections prepared for both Akrofu and Green Valley rock samples

were viewed under the petrographic microscope in both PPL and XPL. The results were recorded in Tables 1 and 2.



Figure 4: Megascopic descriptions of rocks of coarse-grained minerals of feldspars, quartz, biotite and hornblende sampled at: (A) Green Valley; and (B) Akrofu in Volta Region of Ghana.

Table 1: Modal Composition of the Biotite Granitic Gneiss from Green Valley		
Mode	Volume (%)	Characteristics
Quartz	30	Medium grained, subhedral to anhedral habit, recrystallization of quartz
Orthoclase	25	Subhedral to euhedral habit, very coarse-grained, one cleavage plane
Plagioclase	20	Subhedral to euhedral habit, one cleavage plane, shows lamellar twinning. Some are recrystallized
Biotite	10	Irregular flakes and elongated plates, brown to pale brown in PPL and dark brown colour in XPL (pleochroic)
Amphibole	8	Sub- to euhedral, two cleavages planes, cleavages at 120 and 60 degrees
Muscovite	5	Subhedral to euhedral habit, parallel cleavage
Sericite	2	Alteration from the feldspars

Table 2: Modal Composition of the Granodiorite from Akrofu		
Mode	Volume (%)	Characteristics
Quartz	35	Subhedral to anhedral habit, undulose extinction due to deformation and recrystallization of quartz
Plagioclase	25	Subhedral to euhedral habit, one cleavage plane
Orthoclase	16	Subhedral to euhedral habit, coarse-grained, one cleavage plane
Biotite	10	Irregular flakes and elongated plates, brown to pale brown in PPL and dark brown colour in XPL (pleochroic), the mineral shows some rate of alteration and shearing
Amphibole	5	Sub- to euhedral, two cleavages planes, cleavages at 120 and 60 degrees
Muscovite	7	Sub- to euhedral, two cleavages planes, cleavages at 120 and 60 degrees
Sericite	2	Subhedral to euhedral habit, parallel cleavage Alteration from the feldspars




Figure 5: Photomicrograph of representative sample of: (A) gneiss consisting of phenocrysts of Biotite (Bt), Quartz (Qtz), Hornblende (Hbl) at Green Valley; (B) granodiorite consisting of phenocrysts of Plagioclase (Pl), Quartz (Qtz), Biotite (Bt), in XPL at Akrofu

4.1.3 Strength Properties Data Analysis

The Table 3 below is the result obtained after conducting the Schmidt hammer test for both rock samples from Akrofu and Green Valley.

The strength parameters of the rock are classified using Bieniawski classification of strength of intact rock (Bieniawski, 1998). The unconfined compressive strength tests classification scheme is shown in Table 4 below.

Table 3: Schmidt Hammer Test for Akrofu and Green Valley Rock Samples

SAMPLE ID	REBOUND (H)	UCS (MPa)	SAMPLE ID	REBOUND (H)	UCS (MPa)
AKROFU	22	12.00	GREEN VALLEY	38	40
	21	11.00		36	36
	36	36.00		38	40
	21	11.00		33	33
	36	36.00		34	34
	35	35.00		38	40
	MEAN	23.50		MEAN	37.17

Table 4: Classification of Strength of Intact Rock (Bieniawski, 1998)

Qualitative Description	Compressive Strength (MPa)	Point Load Strength (MPa)	Ratings
Extremely Strong	>250	8	15
Very Strong	100 – 200	4 – 8	12
Strong	50 – 100	2 – 4	7
Medium Strong	25 – 50	1 – 2	4
Weak	5 – 25	Use of UCS preferred	2
Very Weak	1 – 5	-do-	1
Extremely Weak	< 1	-do-	Q

5. DISCUSSION OF RESULTS

5.1 Macroscopic Discussion on Both Green Valley and Akrofu Rock Samples

The Green Valley rock sample is grey in colour, coarse-grained metamorphosed rock showing compositional banding with a clear cleavage. The minerals appear in bands with the darker bands having more mafic minerals such as biotite (those containing more magnesium and iron) and the lighter bands having more felsic minerals such as quartz, muscovite and feldspar (containing more of sodium and potassium). The rock has distinct banding due to the appearance of the darker minerals in bands and is distinguished from other rocks such as schist by a well-developed gneissosity. The Akrofu rock sample on the other hand, is medium to coarse-grained rock that is among the most abundant intrusive igneous rocks. It contains quartz and is distinguished from granite by having more plagioclase feldspar than orthoclase feldspar. Hornblende, biotite, quartz and minor amount of muscovite mica are minerals observed in the rock sample. This rock is intrusive igneous rock that have phaneritic texture (i.e. the crystals are so coarse that individual minerals can be distinguished with the naked eye). The grain sizes are visible to the naked eye and its formation is as a result of slow cooling crystallization below Earth's surface.

5.2 Microscopic Discussion on Both Akrofu and Green Valley Rock Samples

The rocks are different from each other but they both show some similar mineralogical content in that minerals like biotite, quartz and hornblende are present in both Akrofu and Green Valley rocks. The Akrofu rocks are characterized by dark biotite minerals arranged in bands giving the rock a coarse texture. The rock has minor content of muscovite giving it a light grey colour and mostly composed of feldspars and quartz. The quartz has hypidiomorphic texture which means some grains of some mineral species are anhedral to euhedral. The potash feldspars are present as subhedral to anhedral microcline crystals and occur as small interstitial crystals or enclosed in plagioclases. The K-feldspar undergoes an altered twinning. Hornblende grains are intertwined with the plagioclase. Crystals are evenly distributed and slightly fractured. Deformation features and kinematic indicators are observed on the Akrofu rock under the microscope and these are the fractures within the grains. The Akrofu granitoid is said to be a granodiorite with regards to its mineralogical composition (Table 2, Figure 5B).

The Green Valley rock sample shows assemblage of hornblende, biotite, quartz, plagioclase feldspar minerals as well as brown to grey phenocrysts of K-feldspars. The quartz has a well-developed distinctive crystal shape with ophitic texture and undergoes undulose extinction. Plagioclase undergoes twinning with tabular crystals. It displays striations and crystals are in a grouping of matrix. Hornblende has an elongated crystals and excellent cleavage. The rock presents foliations which are formed

when the flat or elongate minerals are squeezed by pressure to become aligned and these foliations are highlighted by alternating biotite aligned in a preferential order. The rock does not preferentially break along the plane of foliation because less than 50 % of the minerals formed during the metamorphism are aligned in thin layers. The presence of foliation in the rock is as a result of the re-alignment of minerals when they were subjected to high temperatures and pressures. The Green Valley gneiss is said to be a biotite gneiss because it contains about 10 % of the rock's mineral composition.

5.3 Discussion on the Strength Properties of Both Akrofu and Green Valley Rocks

From Table 3, the rock sample from Akrofu has a compressive strength of 23.50 MPa which falls within the range of 5 – 25 MPa according to the strength classification of intact rocks (Bieniawski, 1998). The rocks that fall in the range 5 – 25 MPa are classified as weak rocks. From Table 3, the rock sample from Green Valley has a compressive strength of 37.17 MPa which falls within the range of 25 – 50 MPa according to the strength classification of intact rocks (Bieniawski, 1998). The rocks that fall in the range 25 – 50 MPa are classified as medium strong rocks. The rocks under study (i.e. Akrofu and Green valley rocks) failed, when subjected to loads beyond their strength and the failure in the rocks took place under compressive, tension and shear forces at different values. It is however, the unconfined compressive strength, which is taken as the most important index property of rocks. The low values obtained for compressive strength are direct effects of the humid tropical conditions that causes weathering of the rocks at the base of the overburden aided by water.

In addition to weathering, structural defects such as fractures, joints, faults and the mechanical drilling process may tend to introduce micro-pores or fractures in the samples thereby leading to a decrease in the strength and modulus of these rocks. The rocks are extensively used in materials for construction. They are characterized by very high crushing (compressive) strength and hence can be easily trusted in most of construction works. The Green Valley rock does not split along the plane of weakness like most other metamorphic rocks and this allow contractors to use it as a crushed stone in road construction. They can be used as building material such as flooring, facing stones on building and also as an architectural stone. On the other hand, the Akrofu rock sample is typically impervious and hard, and forms very strong foundations for most civil engineering works. They are often used as crush stones for road building, construction material, building facade and paving, and as an ornamental stone.

6. CONCLUSION

The Dahomeyan rocks from Akrofu and Green Valley are classified to be granodiorite and biotite granitic gneiss respectively. From the petrographic studies, granodiorite was characterized by dark biotite minerals arranged in bands giving the rock its foliation and a coarse

texture. The rock contains quartz and is distinguished from granite by it having more plagioclase feldspar than orthoclase feldspar. Hornblende, biotite, quartz and minor amount of muscovite mica are minerals observed in the rock sample. It has minor content of muscovite giving it a light grey color and mostly composed of feldspars, and quartz. On the other hand, the gneiss assemblage include hornblende, biotite, quartz, plagioclase feldspar minerals as well as brown to grey coarse-grained phenocrysts of K-feldspars. The structures observed within the rock sample are fractures and foliations.

The Green Valley contains biotite aligned in a preferential order which indicates the presence of foliations. The foliations are highlighted by alternating dark mafic mineral (biotite), clear quartz and feldspar bands and the presence of foliation indicates that the rock has undergone some form of metamorphism. On the other hand, the structures within the grains of the rock from Akrofu are said to be fractured and these are as a result of weathering processes and climatological changes. The strength properties of the rocks were based on the classification of Bieniawski, 1998. The Akrofu rock is classified to be a weak rock with compressive strength of 23.50 MPa while the Green Valley is classified to be a medium strengthened rock with a compressive strength of 37.17 MPa but these low values were as a result of presence of fractures running through the samples as well as weathering process.

ACKNOWLEDGEMENT

I hereby acknowledge the tireless effort and effective contribution of my co-author towards the production of this article.

REFERENCES

- Affaton, P., Rahaman, M.A., Trompette, R., Sougy, J., 1991. The Dahomeyide orogeny: Tectonothermal evolution and relationships with the Volta basin. In: Dallmeyer, R.D., Lecorché, J.P. (Eds.), *The West African Orogens and Circum-atlantic Correlations*. Springer Verlag, Berlin, Pp. 107–122.
- Agbossoumonde, Y., Menot, R.P., Paquette, J.L., Guillot, S., Yessoufou, S., Perrache, C., 2007. Petrological and chronological constraints on the origin of the Palimé-Amlamé granitoids (South Togo, West Africa): A segment of the West African Craton Paleoproterozoic margin

- reactivated during the Pan-African collision. *Gondwana Research*, Elsevier, 12 (4), Pp. 476-488. (10.1016/j.gr.2007.01.004). (insu-00256370)
- Attoh, K., Dallymeyer, R.D., Affaton, P., 1997. Chronology of nappe assembly in the Pan-African Dahomeyide Orogen, West Africa. *Precambrian Research*, 82, Pp. 135-171.
- Bieniawski, Z.T., 1998. Mechanism of brittle fracture of rock, parts I, II, III. *Int. J. Rock Mech. Min. Sci.*, 4 (4), Pp. 395-430.
- Caby, R., 1989. The Pan-African Belt of West Africa from the Sahara to the Gulf of Benin. In: Rodgers, J., Schaer, J.P.E. (Eds.), *The Anatomy of Mountains Range*. Princeton University Press, Pp. 129–170.
- Dzathor, P.K., 1998. *The Ewe Nation and SASADU: A Brief History*. Accra: Berkadams Limited.
- Ganade de Araujo, C.E., Rubatto, D., Hermn, J., Cordani, U.G., Caby, R., Basei, M.A.S., Weinberg, R.F., Sato, K., 2016. Tightening-up NE Brazil and NW Africa connections: advances in zircon geochronology towards a complete plate tectonic cycle in the Dahomeyide belt of the West Gondwana orogen in Togo and Benin. *Precambrian. Res.*, 276, Pp. 29-42.
- Goudie, A.S., 2006. The Schmidt Hammer in Geomorphological research. *Progress in Physical Geography: Earth and Environment*, 30, Pp. 703-718.
- Holm, F.R., 1974. Petrology of alkali gneiss in the Dahomeyan of Ghana. *Geological Society of America Bulletin*, 85, Pp. 1441-1448.
- Kennedy, W., 1964. The structural differentiation of Africa in the Pan-Africa tectonic episode. UK: Leeds University Institute for African Geology.
- Nude, P.M., Shervais, J., Attoh, K., Vetter, S.K., and Barton, C., 2009. Petrology and geochemistry of nepheline syenite and related carbonate-rich rocks in the Pan- African Dahomeyide orogen, southeastern Ghana, West Africa. *Journal of African Earth Sciences*, 55, Pp. 147-157.

